

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/180374

INTERNATIONAL APPLICATION NO.
PCT/EP97/02597INTERNATIONAL FILING DATE
12 MAY 1997PRIORITY DATE CLAIMED
10 MAY 1996TITLE OF INVENTION
LIQUID FATTY COMPONENT CONTAINING COMPOSITIONAPPLICANT(S) FOR DO/EO/US
RITTER, Heike et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)). FILED IN ENGLISH
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

A SUBMISSION OF ARTICLE 34 AMENDMENTS, ALONG WITH TWO AMENDED PAGES, IS ALSO BEING FILED CONCURRENTLY HEREWITH.

17. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):

Search Report has been prepared by the EPO or JPO \$ 930.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
..... \$ 720.00No international preliminary examination fee paid to USPTO (37 CFR 1.482)
but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$ 790.00Neither international preliminary examination fee (37 CFR 1.482) nor
international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$ 1070.00International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(2)-(4) \$ 98.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$ 930.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$ 130.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	13 - 20 =	0	X \$ 22.00
Independent claims	1 - 3 =	0	X \$ 82.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$ 270.00

\$ 0.00

\$ 0.00

\$

TOTAL OF ABOVE CALCULATIONS =

\$ 1,060.00

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

\$

SUBTOTAL =

\$ 1,060.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

+

TOTAL NATIONAL FEE =

\$ 1,060.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

\$

+

TOTAL FEES ENCLOSED =

\$ 1,060.00

Amount to be:
refunded

\$

charged

\$

a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.b. ☒ Please charge my Deposit Account No. 12-1155 in the amount of \$ 1,060.00 to cover the above fees.
A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 12-1155. A duplicate copy of this sheet is enclosed.NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR
1.137(a) or (b)) must be filed and granted to restore the application to pending status.SEND ALL CORRESPONDENCE TO:
LAW OFFICESUNILEVER UNITED STATES
c/o Mrs. Linda Horvath
Patent Department
45 River Road
Edgewater, New Jersey 07020

FILED: 09 NOVEMBER 1998

AJZ:la1

Anthony J. Zelano
SIGNATURE. 12-30595

Anthony J. Zelano

NAME Representative Capacity
Only 27,969

REGISTRATION NUMBER

09/180374

300 Rec'd PCT/PTO 09 NOV 1998

IN THE UNITED STATES DESIGNATED/ELECTED OFFICE

International Application No. *Yal* : PCT/EP97/02597
International Filing Date : 12 MAY 1997
Priority Date(s) Claimed : 10 MAY 1996
Applicant(s) (DO/EO/US) : RITTER, Heike et al.

Title: LIQUID FATTY COMPONENT CONTAINING COMPOSITION

SUBMISSION OF ARTICLE 34 AMENDMENTS

BOX PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

SIR:

The above-identified International Application was amended under Article 34 of the PCT.
Enclosed are two amended pages which are attached as Annexes to the International Preliminary Examination Report.

However, it is requested that examination in the U.S. National Phase be based on the original PCT application as filed.

Respectfully submitted,

Anthony J. Zelano

by [Signature] Reg no 30515

Anthony J. Zelano (Reg. No. 27,969)
Representative Capacity
MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
Arlington Courthouse Plaza I
2200 Clarendon Boulevard, Suite 1400
Arlington, Virginia 22201
Direct Dial: (703) 812-5310
Facsimile No.: (703) 243-6410
E-Mail Address: zelano@mwzb.com

AJZ:/al
Filed: 9 November 1998

F 7340 (V) LEVER 600X

662113-160374

IN THE UNITED STATES DESIGNATED/ELECTED OFFICE

International Application No. *hal* : PCT/EP97/02597
International Filing Date : 12 MAY 1997
Priority Date(s) Claimed : 10 MAY 1996
Applicant(s) (DO/EO/US) : RITTER, Heike et al.

Title: LIQUID FATTY COMPONENT CONTAINING COMPOSITION

PRELIMINARY AMENDMENT

BOX PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

Prior to calculating the national fee, and prior to examination in the National Phase of the above-identified International application, please amend this application as follows:

IN THE CLAIMS:

Please amend claims 3-6 and 8-13 as follows:

- Claim 3, line 1:** Please delete "any one of claims 1 or 2", and insert --claim 1--.
- Claim 4, line 1:** Please delete "any one of claims 1 to 2", and insert --claim 1--.
- Claim 5, lines 1 and 2:** Please delete "any one of the preceding claims", and insert --claim 1--.
- Claim 6, lines 1 and 2:** Please delete "any one of the preceding claims", and insert --claim 1--.
- Claim 8, line 1:** Please delete "any of the preceding claims", and insert --claim 1--.
- Claim 9, lines 1 and 2:** Please delete "any one of the preceding claims", and insert --claim 1--.
- Claim 10, lines 1 and 2:** Please delete "any one of the preceding claims", and insert --claim 1--.
- Claim 11, lines 1 and 2:** Please delete "any one of the preceding claims", and insert --claim 1--.

662740-120374

Claim 12, lines 1 and 2: Please delete "any one of the preceding claims", and insert

--claim 1--.

REMARKS

The principal purpose of this Preliminary Amendment is to eliminate multiple dependencies in order to avoid extra fees.

Respectfully submitted,

Anthony J. Zelano by [Signature] Reg. No. 30595

Anthony J. Zelano (Reg. No. 27,969)

Representative Capacity

MILLEN, WHITE, ZELANO & BRANIGAN, P.C.

Arlington Courthouse Plaza I

2200 Clarendon Boulevard, Suite 1400

Arlington, Virginia 22201

Direct Dial: (703) 812-5310

Facsimile No.: (703) 243-6410

E-Mail Address: zelano@mwzb.com

AJZ:/lal

Filed: 9 November 1998

F 7340 (V) LEVER 600X

300 Rec'd PCT/PTO 09 NOV 1998

Liquid fatty component containing composition

The present invention concerns an organogel containing composition, the organogel being largely composed of a
5 liquid fatty component and a mixture of sterols.

The organogel of the invention has a firmness larger than that of the liquid fatty component when compared at the same temperature, normally room temperature.

10 The liquid fatty component is herein referred to as liquid fat.

Fats are commonly applied in a wide range of consumer products, including food and cosmetic products, and also in technical, non-consumer areas.

15 In many of such applications, it is desired that the product comprising the liquid fat has some structure or firmness, meaning that the product is not as liquid and pourable as the liquid fat itself when comparing at similar conditions. Several methods are used for providing
20 structure, the use of solid fats in addition to the liquid fat being one of the most commonly applied. In applications where in addition to the fat, also a non-fatty liquid, such as water, is present, also emulsifiers and/or thickeners and/or gelling agents are applied, so that the non-fatty
25 liquid adds significantly to the firmness of the end product. In most applications, and in particular in compositions comprising high amounts of fat, solid fat is applied for structuring (or giving firmness to a products) for technical and practical reasons, in particular in fat
30 continuous products comprising > 50% of fat, preferably >60% of fat. At lower fat levels, a combination of solid fat, emulsifiers and thickeners and/or gelling agents is often used. However, in consumer products, in particular in food products, there is an increased desire to reduce or
35 even do without the presence of solid fats for various reasons. For example, such reasons can be based on health concerns, as were fat is consumed, the consumer desires

products with saturated fatty acid (SAFA) levels as low as possible, and, preferably, with low trans fatty acid levels (e.g., less than 4%). More preferred, fat containing products are even free of any trans fatty acid levels.

- 5 Drawbacks sometimes have to do with increase of costs, and/or negative consumer perception of the ingredients used.

Thus, in one embodiment of the present invention, a method
10 is found to replace part or all of the solid fat in a fat composition composed of solid fat alone or of a mixture of solid and liquid fat, by the use of an organogel as claimed herein. Alternatively, the invention is suitable for the reduction of the structuring material necessary to give
15 firmness to the liquid fat containing composition. Another object of the present invention is to provide a new method for giving firmness to a liquid fat, in particular to a glyceride, without chemical modification.

In the present application, a liquid fat means that the
20 liquid fat is pourable at the temperature its application is envisaged. For most products, and in a preferred embodiment, this means at room temperature (20°C).

Examples of such liquid fats include liquid paraffins and liquid, organic fatty components as often applied in
25 consumer products, for example polyol esters like mono-, di- and triglycerides.

The present invention now provides a composition comprising an organogel, the organogel being largely composed of a
30 liquid fatty component, at least one sterol and at least one sterol ester. It is desired that both are present in an amount of at least 1 wt% each, based on the total amount of liquid fat used in the organogel.

Preferably, all liquid fat present in the composition of
35 this invention is formed into an organogel. In a particular embodiment the organogel comprises at least 75% liquid fat. In one embodiment of this invention, the composition

claimed consists of an organogel composed of a liquid fat, a sterol (or mixture of different sterols) and a sterol ester (or mixture of different sterol esters). In another embodiment, other components are present as well.

5 Preferably, the composition does not comprise water.

Accordingly, the organogel of the present invention can be used in combination with other component, such as solid fat, with water, or combinations thereof, and may comprise any other component commonly applied and depending on the
10 end use desired. Preferably, no components are used which negatively affect the firmness of the organogel itself. It is further preferred that in compositions in which water is present and for which a firmness of the organogel is still desired after a prolonged storage time, additional measures
15 are taken. As an example, it is desired to used components which reduce the water activity.

Thus, in the present composition, still solid fat can be present. For example, in a prior art composition composed of liquid and solid fat, the amount of solid fat can be
20 reduced by the use of a mixture of sterol composed as envisaged in the present invention. In this embodiment it is preferred that not only part or all solid fat is replaced by liquid fat, but that all liquid fat in the composition is present in the form of an organogel. In a
25 highly advantageous embodiment, the composition consists of an organogel and solid fat. It is preferred to use solid fat having a low SAFA level, say less than 10%, and a trans fatty acid level of less than 4%, preferably the solid fat being virtually free of trans fatty acids.

30 As indicated above, in the composition of one of the embodiments of the present invention, other components commonly applied in the composition envisaged can be present. For example, in a margarine type of product, all or part of the solid fat present can be replaced by the
35 organogel, and in addition, other components commonly present in such margarine type products such as solid fat, water, flavours, salt, and the like, can be present.

- In this specification, a sterol is a polycyclic alcohol, with at least 24 carbon atoms and at least 4 condensed rings, with a ring size of at least 3 atoms, preferably a ring size in the range of 3-6 atoms. In a preferred embodiment, the rings are almost planar ring systems such as found in cholesterol. In a further preferred embodiment, the hydroxyl groups (or C-O bonds) are positioned in the same plane as the ring system, i.e. equatorial and not axial.
- Sterol esters are, in this specification, esters of phenolic acids of the sterols defined above. The term phenolic acids relates to the family of cinnamic acids, of which caffeic acid and ferulic acid are examples.
- Very good results were obtained when the esterified sterol has a structure highly similar to that of the free sterol applied. In one particularly preferred embodiment, the free sterol is a ferulic acid, and the phenolic acid esterified sterol is a ferulic acid sterol ester. At present, it is believed that the free sterol or a component of similar chemical structure, and the phenolic acid esterified sterol or a component of similar chemical structure, tend to form aggregates when dissolved in the liquid and that these aggregates show a certain level of network formation or even polymerization in the liquid, which is then reflected in a structuring of the liquid. Hence the structuring of the liquid would resemble the formation of a gel known from aqueous gels. However, applicant does not wish to be bound by this theory.
- Examples of suitable combination of sterols and sterol esters selected from the group of phytosterols are oryzanol and sitosterol (often denoted by β -sitosterol). Also cholesterol is found to be a suitable component that can provide structure to a liquid fatty component when applied at specific levels in combination with other phytosterols.

An additional advantage of the present invention is found

in that most of the sterols applicable for providing structure according to the present invention are components obtainable from natural sources. In a preferred embodiment of the invention, the sterols and/or sterol esters applied
5 are components which can also be found in nature.

For example, oryzanol and sitosterol are present as minor components in many plants. In quite some cases, these are even present in the plants from which triglycerides are obtained. It is to be noted, however, that the sterol
10 (ester) components are not present in amounts sufficient to provide structuring in these natural sources, nor are these present in the mole ratios needed for obtaining sufficient structure. Also, in many cases, not all sterols needed for the structuring are present in the plants from which the
15 oils are obtained. At present, these minor components are often partially removed during oil refining.

Hence, one of the objects of the present inventions is to provide a new method for giving firmness to a liquid fat, in particular to an edible glyceride, without chemical
20 modification of any of the ingredients being needed.

In particular, sterols and sterol esters found to be highly suitable for providing hardness to the liquid are selected from the group of phytosterols. In this invention the term
25 phytosterol is used to cover the whole group of free phytosterols, phytosterol fatty acid esters and (acylated) phytosterol glucosides.

There are three major phytosterols, namely beta-sitosterol, stigmasterol and campesterol. Schematic drawings of the
30 components meant are as given in "Influence of Processing on Sterols of Edible Vegetable Oils", S.P. Kochhar; Prog. Lipid Res. 22: pp. 161-188.

Sitosterol can, for example, be obtained from wood and from refining vegetable oil, and normally comprises also a minor
35 amount of other sterols, like campesterol, stigmasterol, various avenasterols etc. For the present invention, it is not needed that the sterols and/or sterol esters used are

highly pure; some impurities can be present, it is considered not to be of concern in particular, when the polarity is relatively low.

- 5 Oryzanol consist of a mixture of ferulic acid esters of unsaturated triterpene alcohols and is also referred to as gamma-oryzanol. In this invention only the term oryzanol is used. For a further description and schematic drawing of oryzanol, reference is made to "Separation of Vitamin E and
10 gamma-Oryzanols from Rice Bran by Normal-Phase Chromatography", M. Diack and M. Saska, JAOCS Vol. 71, no. 11, pp. 1211. Oryzanol can, for example, be obtained from ricebran, and comprises ferulic acid esters of several phytosterols.
- 15 Cholesterol is, for obvious health reasons, less desired when the use of in food products is envisaged. For any other of the applications, however, it may be very well applicable.
- A highly suitable combination, in particular for food
20 products, was found in the use of both oryzanol and sitosterol.

- The organogel of the invention has a firmness larger than that of the liquid fat when compared at the same
25 temperature, normally room temperature.
- In a highly preferred embodiment, firmness of the organogel consisting of liquid oil and a mixture of sterol and sterol esters according to the invention has a Stevens 4.4 hardness ("Stevens 4.4" value) of at least 20 grams
30 measured at 20°C. The Stevens hardness St, expressed in grams, is determined 1 week after manufacturing the organogel when stored at 5°C and thereafter equilibrated for 24 hours at the temperature as indicated, using a 4.4. mm. diameter cylinder in a Stevens -LFRA Texture Analyzer
35 (ex. Stevens Advanced Weighing Systems, UK) load range 1000 g operated "normal" and set at 10 mm penetration depth and 2.0 mm/s penetration rate.

Thus, Stevens hardness is for this purpose considered a parameter sufficient to distinguish between the liquid fat and the "structured" liquid fat (the organogel).

- 5 The mixture of the sterol(s) and sterol ester(s) of the invention are capable of structuring a liquid fat when added to a liquid fat in an amount of at least 1 wt% each, based on the amount of liquid fat. The sterols and sterol esters have a limited solubility in liquid fat, the
- 10 saturation point depending on the specific components used. In most cases, the amount sufficient for structuring is the level whereby all structure providing sterols and sterol esters are added at a total level equal or preferably, above the saturation point of the solution.
- 15 In particular, the minimum total amount of sterols (i.e. sterols + sterol esters) is at least 2 wt% and preferably 3 wt%, based on the total amount of liquid fat present in the composition.

The sterols and sterol esters are dissolved in the liquid

20 fat, and it is believed, without wishing to be bound by theory, that the structuring is obtained by the sterols and sterol esters which are added at a level above the saturation of the solution, meaning that these are the not-dissolved sterols which are believed to provide the

25 structure to the liquid.

It was found that the molar ratio of sterols and sterol esters should be in the range of 1:10-10:1, preferably 1:5-5:1, whereby a preference exist for 1:3-3:1, towards almost equal molar ratio's, as it was found that an optimal

30 structuring effect was obtained for providing firmness to a liquid fat, in particular a liquid triglyceride.

In one particular and preferred embodiment of the invention, the liquid fat is overall apolar. Surprisingly,

35 it was found that almost all apolar liquid fats can be given firmness by the use of a combination of a sterol (mixture) and a sterol ester (mixture), which preferably is

the ester of the specific sterol (mixture) used.

The invention is in particular advantageous in the foods area, so that the use of edible liquid fat and edible sterol(s) and edible sterol ester(s) is preferred.

- 5 Thus, for food products, the liquid fat is an edible fat, and the sterols used is a mixture of phytosterols, preferably oryzanol and sitosterol at a minimum total weight level of 2%, preferably 4%, with a clear optimum at a molar ratio between 3:1 and 1:3, further preferred
- 10 between 1:2 and 2:1. After dissolution of the sterols in the fatty compound at elevated temperature, improvement of structuring capacity of the sterols was found by rapid cooling.
- 15 In a further preferred embodiment for food products, it was found that sitosterol and oryzanol provide significant and sufficient structure to an edible liquid acyl glyceride when both are present in a total amount of at least 2 wt% each, based on the liquid acyl glyceride fat. An optimum in
- 20 providing hardness to a product was found in the use of oryzanol and sitosterol in about equal molar ratio in combination with the use of a liquid polyol ester, preferably a mono-, di- and/or triacyl glyceride. In addition thereto, the use of sitosterol and oryzanol in
- 25 food products are nowadays believed to add beneficial health effects upon their daily intake, in particular with respect to a reduction of coronary heart diseases. For many applications, and in particular for food, the use of edible liquid acyl glycerides, in particular triacyl
- 30 glycerides is preferred as the liquid fatty component. As suitable triacyl glycerides, fats and oils (here used as synonyms) both vegetable and animal fats obtained from natural and synthetic origin can be used. Preferred are liquid vegetable and/or animal fats obtained from natural
- 35 sources. It is further preferred that these oils comprise a substantial amount of C14 to C22 glycerides, and further preferred a substantial amount of glycerides having C16-C20

triglycerides. The C-numbers refer to the number of carbon atoms per fatty acid group

In another preferred embodiment, the liquid fatty compound
5 comprises a substantial amount of saturated acyl
glycerides, in particular saturated triglycerides, as the
use thereof was found to be beneficial with respect to the
hardness of the product. It was observed that a further
increased amount of the structuring sterols is desired for
10 compositions comprising high levels of unsaturated acyl
glycerides compared to the levels needed for the same
degree of structuring. The less unsaturated glycerides were
used, the harder the product with a same amount of sterols
present. In food products, it may be desired for health
15 reasons, however, to use less saturated glycerides and a
higher amount of sterols.

The vegetable triglycerides often used in consumer products
include those obtained from seeds, beans, fruits and nuts,
or parts of these plant materials such as their germs, and
20 are often obtained by mechanical expelling and/or solvent
extraction. Examples of liquid triglycerides which are in
particular suitable for use in the present invention are
sunflower oil, rapeseed oil, flax or linseed oil, soybean
oil, maizegerm or corn oil, wheatgerm oil, ricebran oil,
25 palm oil, olive oil peanut oil, and the like.

Also, oils of animal origin can be used in the present
invention, and include those obtained from processing fish,
for example, fish oil.

Other liquid fats that can be applied in the present
30 invention comprise or consist of sucrose polyol polyesters.

Within the scope of the present invention is a composition
according to the main claim, in which the fat applied is a
fat which is not liquid at room temperature. The term
35 liquid fat regards fat which is liquid at the temperature
applied. For example, fats can be applied at temperatures
higher than room temperature whereby it is still desired

that the fat at its application temperature is not liquid, but has some structure. As is well known, most fats melt at their heating, and also above their melting temperature they can be structured by means of the sterols as presently envisaged within the present invention. It will still be desired to use an amount at or above the level of saturation at the envisaged application temperature. Preferably, temperature of application of the organogel containing composition is less than 80° C, as it was found that the organogel becomes less stable at temperatures higher than that.

In another embodiment of the invention, a liquid fat or oil of vegetable or animal source is used as indicated above, to which, in addition to the sterols, at least 1 wt% based on the amount of liquid fatty compound used, of monoglycerides is added. It was found that an additional increase of hardness is obtained by the addition of 1% or more, e.g. 2-10% of monoglyceride, or that a reduction of the amount of sterols needed to obtain hardness or viscosity for the liquid fatty compound by the addition of a monoglyceride.

The composition according to the present invention can be prepared by the simple dissolution of the sterol(s) in the liquid fat. For example, a composition of significant hardness compared to the liquid itself is obtained by dissolving a phenolic acid ester of a sterol and a sterol by mixing the ingredients and stirring at elevated temperatures.

For many fats and oils, a temperature of more than e.g. 40-50°C, often more than 75°C or 85°C will be sufficient, in some cases 90°C degrees is desired in order to achieve rapid dissolving. As soon as a clear, transparent solution is obtained the liquid solution can be allowed to cool to, for example, ambient temperature.

The formation of the structured system often proceeds

rather slowly, and for some cases, it can take more than one day up to several days before the ultimate degree of structuring is achieved. It was found that a more rapid and even instantaneous structuring of the solution is obtained
5 by applying rapid cooling. A very suitable means therefore is a surface scraped cooling device, for example a votator (A-unit).

From a nutritional point of view, it is desired that in
10 food products as little saturated fat (saturated fatty acid groups containing triglycerides) are present. Therefore, in these days, it is often desired to use untreated oils and fats having a high amount of unsaturated fatty acid groups. Such oils, however, are very often pourable at room
15 temperature and therefore less suited for use in products that should have some firmness, such as spreads, dressings, mayonnaise and even squeezable margarines used as vegetable topping, creams, fillings and toppings. Up till now, these oils were treated by which treatment some firmness or
20 hardness was obtained. However, such treatment has the serious disadvantage of saturating the unsaturated fatty acid groups. It is desired from a nutritional point of view, therefore, to use as much untreated oils and fat in a food product as the hardness allows to. With the present
25 invention, now, firmness to a liquid oil or fat can be given by the addition of at least one sterol and at least one sterol ester in an amount of at least 1 wt% each, based on the liquid fat to which firmness is to be given.

30 The organogel is preferably used in consumer goods, such as cosmetic products or food products. Also these products comprising such a composition are part of the invention. Thus, the present invention allows to prepare fat based food products in which a dispersion of an organogel is
35 used. Although it is very well possible to prepare food products in which all fat is present in the form of an organogel according to the present invention, in one

embodiment the food product comprises fat in the form of a organogel as defined in this application and another (solid) fat.

5 The present invention is now further illustrated by way of following examples.

In the examples, percentages are weight percentages, unless indicated otherwise.

The degree of structuring of the samples was determined by
10 means of a Stevens penetrometer under the conditions:

- probediameter: 6.4 mm;
- penetration depth: 20 mm;
- penetration rate: 20 mm/s;
- temperatures: ambient (ca 20°C).

Example I

For the determination of the solubility of the ricebran specific phytosterol "oryzanol", comprising ferulic acid esters for several phytosterols, in a refined sunflower oil, 1-10 % of the pure oryzanol (obtained from Tsuno Rice Fine Chemicals Co Ltd, Japan, > 98% ferulic acid esters) and β -sitosterol (ex Kaukas Oy, Chemical Mill, Finland, comprising 92% B-sitosterol, 7% campesterol and about 1% artenols. Average Mw was 413), were admixed to the oil at elevated temperatures between 50 and 100° C, while stirring until a clear and transparent solution was obtained.

The solubility of oryzanol in sunflower oil, defined as the (w/w) concentration that gives a clear solution after a prolonged keeping time at a certain temperature, was established to increase about linearly from ca 2 % at 20 °C to ca 10 % at 90 °C.

When a saturated solution was allowed to cool down and kept for a prolonged time in order to cope with often occurring retarded crystallization - supersaturation - the solution showed the formation of discrete oryzanol crystals at the bottom of the jar. Under these circumstances no structuring of the oil phase was observed at all (= experiment no. I.1).

Similar solubility determinations were performed with a pure wood based sitosterol product consisting of ca 92 % of genuine sitosterol and about 8 % of campesterol.

The solubility of this sitosterol in refined sunflower oil was found to be ca 1 % at 20 °C and linearly increasing to ca 4 % at 90 °C, the presence of a small amount of it in the pure oil not taken into account.

When the temperature was allowed to drop to below the saturation temperature the sitosterol solution too showed

precipitation of sitosterol crystals after some time.
Also in these solubility tests no structuring phenomena
were noticed at all (= experiment nr. I.2).

- 5 In order to check the solubility for the different
phytosterols, solubility experiments were also carried out
with combinations of both the sterols dissolved in the oil.
These solutions of the combined phytosterols remained
liquid as long as the temperature was kept above say 50 °C.
- 10 Very surprisingly it was established that, starting at a
level of approx. 2 % of oryzanol and 2 % of sitosterol
dissolved in the sunflower oil, the solution of the
phytosterols became jelly and even solid after some days
- 15 when the solutions were cooled down to ambient temperatures
and stirring was not applied anymore (= experiment no.
I.3).

Example II

- 20 The unexpected structuring capability of the combined
dissolving of oryzanol and sitosterol in liquid sunflower
oil was further elucidated by performing a series of
dissolving experiments with the mass ratio
- 25 oryzanol:sitosterol as a variable.
The constant conditions in this series of experiments
follow from the description of the treatment:
- put 94.0 g of refined (and winterized) sunflower
oil in a stirred glass beaker;
 - 30 - admix the required amounts of oryzanol and
sitosterol to the oil at a constant total sterol
concentration level of 6.0 %;
 - increase the temperature of the mixture to 90 °C
while stirring;
 - 35 - continue stirring for 30 min after achieving a
clear solution;
 - allow the solution to cool to ambient

- temperature;
- keep the solution at rest at ambient temperature for 6 days;

5 The Stevens' hardness results in grams force measured in solutions containing varying amounts of oryzanol and sitosterol, are listed in Table II. Experiment II.1 and II.9 are comparative Examples.

10

TABLE II

15

exp.no:	oryzanol [g]	sitosterol [g]	mol ratio ory:sito	hardness [g]
II.1	5.55	0.45	8.36	0
II.2	5.01	0.99	3.46	13
II.3	4.43	1.57	1.92	25
II.4	3.90	2.10	1.27	37
II.5	3.56	2.44	1.00	57
II.6	3.54	2.46	0.98	57
II.7	2.96	3.04	0.66	81
II.8	2.01	3.99	0.34	31
II.9	0.96	5.04	0.13	0

20

This series of experiments clearly shows the structuring phenomenon when the two phytosterols considered are dissolved in the sunflower oil together.

25

Example III

30

The total phytosterol concentration is illustrated by the

following example:

Ratio and total weight amount are listed in the table below.

- 5 The other constant conditions during these tests were as described in example II, except for a variable amount of sunflower oil involved (listed in the table) and the keeping time, which was 3 days in this experiment. The results of these treatments are listed in the following
- 10 table:

TABLE III

exp. no	oryzanol [g]	sitosterol [g]	oil [g]	sterols [%]	hardness [g]
III.1	2.37	1.63	96.0	4.0	0
III.2	3.56	2.44	94.0	6.0	26
15 III.3	5.34	3.66	91.0	9.0	1057
III.4	7.12	4.88	88.0	12.0	1901
III.5	9.49	6.51	84.0	16.0	3520

- From this table it follows that the hardness of the
- 20 structured sunflower oil increases substantially when increasing the total amount of the phytosterols. Example III.1 turned out not to have any hardness at the time of measurement. However, in an additional experiment, a composition the Stevens hardness was found to be about 20
- 25 after 26 days

Example IV

In this series of experiments the varied conditions were:

- holding time after cooling: 1 - 4 - 7 days;
- 5 - holding temperature: 5 - 15 - 25 °C.

The total sterol concentration was kept at 5 % now and the molecular ratio oryzanol:sitosterol 1:1. The Stevens hardness measurement was performed at ambient temperature. The dissolving was performed in a stirred and baffled,
10 double walled, glass reactor. The results are listed in the following table:

"effect of time, temperature and sterol ratio on hardness"

	exp. no.	time (days)	temp. (°C)	hardness (g) or:sito 1:1	hardness (g) or:sito 2:1	hardness (g) or:sito 1:2
15	IV.1	1	5	0	807	565
	IV.2	1	15	0	857	548
	IV.3	1	25	0	5	14
	IV.4	4	5	185	860	568
	IV.5	4	15	1	753	623
	IV.6	4	25	0	90	38
20	IV.7	7	5	209	920	622
	IV.8	7	15	18	783	661
	IV.9	7	25	10	254	190

From these results it is clear that the structuring
25 requires several days and that it proceeds faster at the lower temperature. Again, at a further prolonged standing of the compositions, the hardness was found to further increase. A rapid structuring occurred at using a votator (A unit) for cooling down the compositions.

30

Example V

Several different types of liquid triglyceride oils, with a different degree of unsaturation were used in this
35 experiment. The oils used were obtained from: palm olein, olive, ricebran, rapeseed, sunflower and flax or linseed oil. The palm olein stemmed from a dry fractionation at 15 °C.

These experiments were performed at the following conditions:

- total sterol concentration: 6 %;
- weight ratio oryzanol:sitosterol: 1:1;
- 5 - dissolving temperature: 90 °C;
- holding time: 6 days;
- holding temperature: 20 °C.

The Stevens' hardness results achieved are listed below:

10

TABLE V

exp. no	oil	hardness [g]
V.1	olive	127
V.2	rice bran	73
15 V.3	rapeseed	84
V.4	sunflower	81
V.5	flax seed	22

Example VI

- 20 The influence of using the animal cholesterol instead of sitosterol on the structuring behaviour was checked by performing structuring with 46 g of sunflower oil + 3 g of oryzanol + 1 g of cholesterol.

- 25 Soon after dissolving the animal and the vegetable sterols in sunflower oil at 90 °C the structuring behaviour became visible. See the next table:

TABLE VI

exp. no	2nd component	description	hardness [g]
VI.1	sitosterol	reference	81
30 VI.2	cholesterol	rapidly solidified	78

During the first couples of days the cholesterol structured oil looked very transparent but after some time the sample became more "hazy" like with all previous tests.

5 Example VII

The refined sunflower oil used for the trials described sofar still can contain small amounts of polar matter like di- and monoglycerides and free fatty acids.

For testing sterols structuring in purified triacylglycerol
10 refined and winterized sunflower oil was additionally SiO₂-column treated at ambient temperature under the following conditions:

1 part of oil dissolved in 2 parts of hexane;

column treatment with silicagel (no 60 ex Merck)

15 In the purified oil 3% of oryzanol and 3% of sitosterol were dissolved at 90°C. After 6 days holding at ambient temperature a very hard structure was obtained with the purified triglyceride oil reflected in 1049 g Stevens' hardness.

20

Example VIII

The influence of the presence of varying amounts of monoglycerides in refined sunflower oil was established by structuring experiments nos. VIII.1-4 without and with

25 admixing 3, 6 or 9% of pure monoacylglycerol (mag) (Hymono)
The hardness results are listed in the table below:

TABLE VIII

exp. no	description	hardness [g]
VIII.1	sunflower ref.	81
VIII.2	3% monoacyl glycerol (MAG)	168
VIII.3	+6 % MAG	325
VIII.4	+9% MAG	441

30

TABLE VIII

exp. no	description	hardness [g]
VIII.1	sunflower ref.	81
VIII.2	3% monoacyl glycerol (MAG)	168
VIII.3	+6 % MAG	325
VIII.4	+9% MAG	441

From this table it follows that the addition of pure monoglycerides has a positive effect on the hardness of the sterols structuring.

Example IX

A series of structuring tests was performed with a number of different organic non-triglyceride liquids with 6 % total sterols concentration and at a molar ratio of 1:1 of oryzanol:sitosterol. The results are given in the following table:

"phytosterol structuring in non-TAG liquids"

exp.nr.	liquid	type	hardness (g)
IX.1	water	inorganic	0
IX.2	methanol	alcohol	0
IX.3	ethanol	-	0
IX.4	isopropyl alcohol	-	0
IX.5	propylene glycol	diol	131
IX.6	glycerol	triol	0
IX.7	acetone	ketone	0
IX.8	acetic acid	acid	0
IX.9	oleic acid	-	< 5
IX.10	hexane	alkane	26
IX.11	paraffine oil	-	867

The results show quite significant structuring ability by phytosterols in paraffine oil, propylene glycol, hexane

and to a certain extent oleic acid.

Example X

5

Mixing sterols structured sunflower oil at ambient temperature with demineralized water at increasing oil:water ratio, in steps of 10%, resulted in white coloured stable soft emulsions from a 50:50 ratio or

10 higher.

At lower ratio's the oil phase remained as a separate phase in the water.

When mixing oryzanol and sitosterol containing sunflower oil - total sterol concentration of about 6% and at a 1:1
15 molar ratio -with 20% of water, a quite hard margarine like spread was obtained by applying rapid cooling in a votator (A-unit) at a temperature profile of 60-90°C to 0-5°C.

After a few days in the fridge, the spread obtained became
20 soft and could very well be applied as a squeezable margarine.

EXAMPLE XI

25

In this experiment, spreads were prepared comprising the organogel and water. The organogel was prepared from ultrasitosterol (a sterol obtained from wood pulp), the oryzanol was obtained from Kaukas Oy.

30

Two series were prepared, both prepared by preparing two separate phases, i.e. an aqueous and a fat phase, and combining these phase after a pre-treatment in a C-unit. The pre-treatment of the fat phase encompassed the heating
35 of the fat phase ingredients to 90°C, cooling to 60°C, passing the composition through an A unit so as to rapidly cool down to 15°C.

RECTIFIED SHEET (RULE 91)
ISA/EP

The aqueous phase was prepared by heating the combined ingredients to 60°C, passing it through an A-unit so as to rapidly cool down to 15°C

- 5 In Example XI.A, the aqueous phase contained, salt, syrup, or no additives; the concentration of sterol:sterolesters (1:1 m/m) was 5 wt% on the fat phase.

Example XI.A encompasses 6 experiments, including 1 reference example. Results are provided in the following

10 table:

TABLE XI.A

exp	fat + sterols [%]	H ₂ O [%]	NaCl [%]	Syrup [%]	Stevens after 2 days	Stevens after 5 days
A.1	84.5	15.5	0	0	70	67
A.2	84.6	12.6	2.77	0	108	125
A.3	82.4	13	4.57	0	100	94
A.4	81.4	7.0	0	11.6	117	122
A.5	81.0	2.7	0	16.4	141	153

- 20 In Example XI.B, the aqueous phase contained about 12 % of the total fat content, and so, the aqueous phase was a pre-mix in itself. All sterols were added to the fat phase. Accordingly, the sterol concentration in the fat phase was 5.65%, whereas the sterol concentration based on the total
- 25 amount of fat in the composition was 5%. To the fat phase, 0.5% Dimodan BP and a minor amount of carotene were added.

TABLE XI.B

exp.	fat+ sterols [%]	H ₂ O [%]	NaCl [%]	Syrup [%]	Stevens after 2 days	Stevens after 5 days
B.1	95.5	4.5	0	0	34	36
B.2	95.5	3.7	0.81	0	133	138
B.3	95.4	3.4	1.19	0	157	192
B.4	95.3	1.8	0	2.97	105	120
B.5	94.8	0.7	0	4.45	197	203
ref	100	0	0	0	245	302

5

10

662740-42830

CLAIMS

- 1 Composition containing an organogel, the organogel comprising a liquid fatty component, at least one sterol and at least one sterol ester.
- 2 Composition according to claim 1, wherein the total amount of sterol and the total amount of sterol ester is at least 1 wt% for each, based on the amount of the liquid fatty component.
- 3 Composition according to any one of claims 1 or 2, wherein composition consists of a liquid fat, a sterol or mixture of different sterols, and a sterol ester or mixture of different sterols, and, optionally, monoglycerides.
- 4 Composition according to any one of claims 1 to 2, wherein the composition does not comprise water.
- 5 Composition according to any one of the preceding claims, wherein the sterol ester has a structure highly similar to that of the sterol applied.
- 6 Composition according to any one of the preceding claims, wherein the sterol and the sterol esters are selected from the group of phytosterols.
- 7 Composition according to claim 6, wherein the sterol is a sitosterol and the sterol ester is oryzanol.
- 8 Composition according to any of the preceding claims, wherein the organogel in the composition has a stevens 4.4. harness of at least 20 grams, measured at 20°C.

RECTIFIED SHEET (RULE 91)
ISA/EP

- 9 Composition according to any one of the preceding claims, wherein the total amount of sterols is at least 3 wt%, based on the total amount of liquid fat present in the composition.
- 10 Composition according to any one of the preceding claims, wherein the molar ratio of sterols and sterol esters is in the range of 1:5 to 5:1, preferably in the range of 1:3 to 3:1, more preferred in an almost equal molar ratio.
- 11 Composition according to any one of the preceding claims, wherein monoglycerides are present.
- 12 Composition according to any one of the preceding claims, wherein the composition is a food product and the fat present is edible fat, and the sterols and sterol esters applied are edible components.
- 13 Composition according to claim 12, wherein the food product is a spread, a squeezable margarine, a dressing or mayonnaise.

CLAIMS

- 1 Composition containing an organogel, the organogel comprising a liquid fatty component, at least one sterol and at least one sterol ester, wherein the molar ratio of sterols to sterolesters is in the range of 1:5 to 5:1.
- 2 Composition according to claim 1, wherein the total amount of sterol and the total amount of sterol ester is at least 1 wt% for each, based on the amount of the liquid fatty component.
- 3 Composition according to any one of claims 1 or 2, wherein the composition consists of a liquid fat, a sterol or a mixture of different sterols, and a sterol ester or a mixture of different sterols, and monoglycerides.
- 4 Composition according to any one of claims 1 to 2, wherein the composition does not comprise water.
- 5 Composition according to any one of the preceding claims, wherein the sterol ester has a structure highly similar to that of the sterol applied.
- 6 Composition according to any one of the preceding claims, wherein the sterol and the sterol esters are selected from the group of phytosterols.
- 7 Composition according to claim 6, wherein the sterol is a sitosterol and the sterol ester is oryzanol.

AMENDED SHEET,

- 8 Composition according to any of the preceding claims, wherein the organogel in the composition has a Stevens 4.4. hardness of at least 20 grams, measured at 20°C.
- 9 Composition according to any one of the preceding claims, wherein the total amount of sterols, ie sterols plus sterolesters, is at least 3 wt%, based on the total amount of liquid fat present in the composition.
- 10 Composition according to any one of the preceding claims, wherein the molar ratio of sterols and sterol esters is in the range of 1:3 to 3:1, more preferred in an almost equal molar ratio.
- 11 Composition according to any one of the preceding claims, wherein monoglycerides are present.
- 12 Composition according to any one of the preceding claims, wherein the composition is a food product and the fat present is edible fat, and the sterols and sterol esters applied are edible components.
- 13 Composition according to claim 12, wherein the food product is a spread, a squeezable margarine, a dressing or a mayonnaise.

AMENDED SHEET

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Includes Reference to PCT International Applications)

Attorney Docket No.
F 7340 (V)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

LIQUID FATTY COMPONENT CONTAINING COMPOSITION

the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application Serial No. _____ on _____ and was amended on _____ (if applicable)

☒ was filed as PCT international application PCT/EP/02597 on 12 May 1997 and was amended under PCT Article 19 on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (if PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 U.S.C. 119
Europe	96201303.3	10 May 1996	Yes

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code §112. I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.


PRIOR U.S. PROVISIONAL APPLICATION(S) FOR BENEFIT UNDER 35 U.S.C. 119(e)

APPLICATION NUMBER	DATE OF FILING (day, month, year)

hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code §112. I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120.

U.S. APPLICATIONS		STATUS (CHECK ONE)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED



COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Includes Reference to PCT International Applications)	Attorney Docket No. F 7340 (V)
--	-----------------------------------

PCT APPLICATIONS DESIGNATING THE U.S.				
PCT APPLICATION NO.	PCT FILING DATE	U.S SERIAL NUMBERS ASSIGNED (if any)		
PCT/EP97/02597	12 May 1997			

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agents(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number)

James J. Farrell, Esq., Reg. 26,162
Gerard J. McGowan, Jr., Esq., Reg. 29,412
Matthew Boxer, Esq., Reg. 28,495

Milton L. Honig, Esq., Reg. 28,617
Ronald A. Koatz, Esq., Reg. 31,774
Neil Y. Gilbert, Esq., Reg. 35,156

A. Kate Huffman, Esq., Reg. 31,372
Rimma Mitelman, Reg. 34,396

Send correspondence to:

Unilever
Patent Department
45 River Road, Edgewater, New Jersey 07020

Direct Telephone Calls to: (name and telephone #):

James J. Farrell
(201) 943-7100 Ext. 2332

201


FULL NAME OF INVENTOR <u>Heide</u>	FAMILY NAME RITTER	FIRST GIVEN NAME Heike	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Kleve	STATE OR FOREIGN COUNTRY Germany <u>DEX</u>	COUNTRY OF CITIZENSHIP German
POST OFFICE ADDRESS	POST OFFICE ADDRESS Kaplan-Mertens Strasse 9	CITY D-47533 Kleve	STATE & ZIP CODE/COUNTRY Germany

202

FULL NAME OF INVENTOR <u>Leo</u>	FAMILY NAME SANDE van de	FIRST GIVEN NAME Robert	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Vlaardingen	STATE OR FOREIGN COUNTRY The Netherlands <u>NLX</u>	COUNTRY OF CITIZENSHIP The Netherlands
POST OFFICE ADDRESS	POST OFFICE ADDRESS UNILEVER RESEARCH VLAARDINGEN, Olivier van Noortlaan 120	CITY 3133 AT Vlaardingen	STATE & ZIP CODE/COUNTRY The Netherlands

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201 Heike Ritter <u>Heide Ritter</u>	SIGNATURE OF INVENTOR 202 Robert Leo K. M. van de Sande <u>Robert van de Sande</u>
DATE <u>November the 24th, 1998</u>	DATE <u>December 2nd, 1998</u>



COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Includes Reference to PCT International Applications)	Attorney Docket No. F 7340 (V)
--	-----------------------------------

203

FULL NAME OF INVENTOR 3-00	FAMILY NAME MULLER	FIRST GIVEN NAME Volkmar	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Vlaardingen	STATE OR FOREIGN COUNTRY The Netherlands NLX	COUNTRY OF CITIZENSHIP The Netherlands
POST OFFICE ADDRESS	POST OFFICE ADDRESS UNILEVER RESEARCH VLAARDINGEN, Olivier van Noortlaan 120	CITY 3133 AT Vlaardingen	STATE & ZIP CODE/COUNTRY The Netherlands

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 203 Volkmar Muller		
DATE December 21st 88		

602740-1203760